

LITERATURA

1. Nicol AL, Adams MCB, Gordon DB, et al. AAAPT diagnostic criteria for acute low back pain with and without lower extremity pain. *Pain Medicine*. 2020;21(11):2661-2675. doi: 10.1093/pm/pnaa239.
2. Macías-Toranzo I, Rojas-Ocaña MJ, Sanchez-Ramos JL, et al. Pain catastrophizing, kinesiophobia and fearavoidance in non-specific work-related low-back pain as predictors of sickness absence. 2020;15(12):e0242994. Available from: <https://doi.org/10.1371/journal.pone.0242994>.
3. Ferguson SA, Merryweather A, Thiese MS, et al. Prevalence of low back pain, seeking medical care, and lost time due to low back pain among manual material handling workers in the United States. *BMC Musculoskelet Disord*. 2019;24:20. Available from: <https://doi.org/10.1186/s12891-019-2594-0>.
4. Sauter M, Barthelme J, Müller C, et al. Annual handling of heavy loads and low back pain among different occupational groups: results of the 2018 BIBB/BAuA employment survey *BMC Musculoskelet Disord*. 2021;22:956. Available from: <https://doi.org/10.1186/s12891-021-04819-z>.
5. Bleecker ML, Hansen JA. *Occupational neurology and clinical neurotoxicology*. Williams and Wilkins, Baltimore, 1994.
6. Newington L, Harris EC, Walker-Bone K. Carpal tunnel syndrome and work *Best Pract Res Clin Rheumatol*. 2015;29(3): 440-453. doi:10.1016/j.berh.2015.04.026.
7. Miettinen L, Ryhänen J, Shiri R, et al. Work-related risk factors for ulnar nerve entrapment in the Northern Finland Birth Cohort of 1966. 2021;11:10010. Available from: <https://doi.org/10.1038/s41598-021-89577-7>.
8. Jackson JA, Olsson D, Burdorf A, et al. Occupational biomechanical risk factors for radial nerve entrapment in a 13-year prospective study among male construction workers. *Occup Environ Med* 2019;76:326-331. doi:10.1136/oemed-2018-105311.
9. Hogg W, Togliati M. Overuse cervical dystonia: A case report and literature review. *Tremor and Other Hyperkinetic Movements*, 2016. Available from: <http://www.tremorjournal.org>.
10. Schiellen L, Tizek L, Biedermann T, et al. Tick bites in different professions and regions: pooled cross-sectional study in the focus area Bavaria, Germany. *BMC Public Health*. 2022;22:234. Available from: <https://doi.org/10.1186/s12889-021-12456-3>.
11. Jozefowicz-Korczynska M, Zamysłowska-Szmytko E, Piekarska A, et al. Vertigo and severe balance instability as symptoms of Lyme disease – literature review and case report. *Front. Neurol*. 2019;10. Available from: <https://doi.org/10.3389/fneur.2019.01172>.
12. Ehler E, Novotná A, Mrklavský M. Poliomyelitis-like syndrome on the background of tick-borne meningoencephalitis. *Cesk Slov Neurol N*. 2007;70/103(4):420-423.
13. Nath A. Neurologic manifestations of severe acute respiratory syndrome coronavirus 2 infections. *Continuum* 2021;27(4):1051-1065.